arXiv:1304.7972v1 [nlin.PS] 30 Apr 2013

Collision of water wave solitons

N. Fenyvesi and G. Bene

Institute of Physics, Loránd Eötvös University
Pázmány Péter sétány 1/A H-1117 Budapest, Hungary

E-mail: nor.fenyvesi@gmail.com, bene@arpad.elte.hu

Abstract. A classification of the time evolution of the two-soliton solutions of the Boussinesq equation is given, based on the number of extrema of the wave. For solitons moving in the same directions, three different scenarions are found, while it is shown that only one of these scenarios exists in case of oppositely moving solitons.

PACS numbers: 47.35.Fg

1. Introduction

Interest in environmental flows[1]-[18] both utilize and motivate theoretical work in nonlinear wave phenomena. Among these are solitons[19]-[26], which appear in many different situations and have applications in several other branches of physics[27], [28]. Although solitons are well known and much studied, we believe that a simple classification scheme of the evolving wave shapes of two colliding water wave solitons may not be in vain.

Weakly nonlinear waves in a shallow channel may be well described in terms of the Boussinesq equation. It allows propagation in both directions.‡ The small parameters of the problem are the ratio of the wave amplitude to the water depth ($\epsilon = a/h$) and the ratio of the water depth to the wave length ($\delta = (h/l)^2$), these are assumed to be comparable. The smaller these parameters are, the better the approximation will be. Therefore, any phenomenon found when solving the Boussinesq equation corresponds to observable effects if ϵ and δ are sufficiently small.

It is well known that the Boussinesq equation is fully integrable and has soliton solutions. Especially, when two solitons have collided, their individual properties are preserved and completely restored when they spatially separate. The change of the wave's shape during collisions is interesting and, as it turns out, follows three possible scenarios. Especially, it is intriguing, what happens if solitons with very similar parameters collide, since it is well known, that the one-soliton solution is unique (up to a scale transformation), hence, two identical, albeit spatially separated solitons do not form an exact solution.

‡ Strictly speaking, the derivation of that equation does not allow counterpropagating waves.

In the present paper we classify the possible wave shape changes during the collision of two Boussinesq solitons.

2. General properties of the two-soliton solution of the Boussinesq equation

The Boussinesq equation in dimensionless form is given by

$$\eta_{tt} - \eta_{xx} - 3\left(\eta^2\right)_{xx} - \eta_{xxxx} = 0 \ . \tag{1}$$

Two-soliton solutions are generated by the Zakharov-Shabat scheme [19], and may be written as

$$\eta = -4\frac{\partial}{\partial x}\left\{\frac{k_1(1+q)+k_2(1+p)-(k_1+k_2)a}{(1+q)(1+p)-a}\right\} \ , \tag{2}$$

where the new variables p and q are defined by

$$p = \mathrm{e}^{2k_1x-2\omega_1t} \ , \quad q = \mathrm{e}^{2k_2x-2\omega_2t} \tag{3}$$

and the parameter a is given by

$$a = \frac{4k_1k_2}{(k_1+k_2)^2+\frac{1}{12}\left(\frac{\omega_1}{k_1}-\frac{\omega_2}{k_2}\right)^2} \ . \tag{4}$$

In these expressions k_1 and k_2 are arbitrary positive parameters characterizing the individual solitons and

$$\omega_j = \pm k_j\sqrt{1+4k_j^2} \ , \quad (j=1,2) \tag{5}$$

with positive sign for a soliton propagating from left to right, and negative otherwise. Note that if $\omega_1 \neq \omega_2$,

$$0 < a < 1 \ . \tag{6}$$

On the other hand, if $\omega_1 = \omega_2$, we also have $k_1 = k_2$ that corresponds to a one-soliton solution, not being of interest here.

An important observation is that in the extreme cases when $k_1,\ k_2 \ll 1/2$ or $k_1,\ k_2 \gg 1/2$ the parameter a depends only on the ratio

$$\kappa = \frac{k_2}{k_1} \ .$$

Explicitly, we get for solitons moving in the same direction

$$a = \begin{cases} \frac{4\kappa}{(1+\kappa)^2+\frac{1}{3}(1-\kappa)^2} & \text{if} \quad k_1, k_2 \gg \frac{1}{2} \\ \frac{4\kappa}{(1+\kappa)^2} & \text{if} \quad k_1, k_2 \ll \frac{1}{2} \end{cases} \tag{7}$$

and for solitons moving oppositely

$$a = \begin{cases} \frac{4\kappa}{(1+\kappa)^2+\frac{1}{3}(1+\kappa)^2} & \text{if} \quad k_1, k_2 \gg \frac{1}{2} \\ 0 & \text{if} \quad k_1, k_2 \ll \frac{1}{2} \end{cases} \tag{8}$$

For the spatial derivative of the wave we have

$$\frac{\partial\eta}{\partial x} = \frac{8}{\left((1+q)(1+p)-a\right)^3}\sum_{n=1}^{5}\sum_{j=0}^{n} c_{nj}p^jq^{n-j} \ , \tag{9}$$

where the coefficients are given by

$$\begin{aligned}
c_{10} &= -k_2^3(a-1)^2 \\
c_{11} &= -k_1^3(a-1)^2 \\
c_{20} &= -k_2^3(a-1) \\
c_{21} &= -(k_1+k_2)\left[a(k_1+k_2)^2 - 3k_1^2 + 3k_1k_2 - 3k_2^2\right](a-1) \\
c_{22} &= -k_1^3(a-1) \\
c_{31} &= (k_1-k_2)\left[a(2k_1^2+5k_1k_2+2k_2^2) - 3k_1^2 - 3k_1k_2 - 3k_2^2\right] \\
c_{32} &= (k_2-k_1)\left[a(2k_1^2+5k_1k_2+2k_2^2) - 3k_1^2 - 3k_1k_2 - 3k_2^2\right] \\
c_{41} &= -k_1^3 \\
c_{42} &= -(k_1+k_2)\left[a(k_1+k_2)^2 - 3k_1^2 + 3k_1k_2 - 3k_2^2\right] \\
c_{43} &= -k_2^3 \\
c_{52} &= k_1^3 \\
c_{53} &= k_2^3
\end{aligned} \tag{10}$$

All the other coefficients are zero. Again, in the extremes $k_1,\ k_2 \ll 1/2$ or $k_1,\ k_2 \gg 1/2$ a common factor k_1^3 can be pulled out of all the coefficients and the rest depends only on the ratio k_2/k_1. Accordingly, for both very small and very large wave numbers the behavior of the solitons is determined by this ratio, up to a scaling.

For minima and maxima of the amplitude $\eta(x,t)$ at a given time t we have

$$\sum_{n=1}^{5}\sum_{j=0}^{n} c_{nj}p^j q^{n-j} = 0\ . \tag{11}$$

Additionally, due to the definitions (5) we have

$$q^{k_1}p^{-k_2} = \mathrm{e}^{2(k_2\omega_1 - k_1\omega_2)t}\ . \tag{12}$$

Thus, extrema are given by the intersections of the curves (11) and (12). The graph of Eq.(12) is a power function with a time-dependent coefficient. In fact, time appears only here. As for the graph of Eq.(11), we may prove some general properties on the basis of Eqs.(10), namely

(i) Exchanging k_1 with k_2 and ω_1 with ω_2 is equivalent with exchanging p with q. This follows directly from Eqs.(2)-(4).

(ii) The curves can have at most three intersections with a line $p = const.$ or $q = const.$ Indeed, for a fixed p Eq.(11) is a third order polynomial of q and vica versa.

(iii) The transformation

$$p \to (1-a)\frac{1}{p}\ , \quad q \to (1-a)\frac{1}{q} \tag{13}$$

leaves Eq.(11) invariant.

Indeed, direct substitution shows that under transformation (13) the expression

$\sum_{n=1}^{5}\sum_{j=0}^{n} c_{nj}p^j q^{n-j}$ goes over into

$$-\frac{(1-a)^3}{p^3q^3}\sum_{n=1}^{5}\sum_{j=0}^{n} c_{nj}p^j q^{n-j}\ . \tag{14}$$

In view of the definition of p and q, the symmetry (13) means that the two-soliton solution is invariant with respect to a simultaneous spatial and temporal reflection, with respect to suitably chosen origins. Explicitly, transformation (13) is equivalent with

$$x \to 2x_0 - x\ , \quad t \to 2t_0 - t\ , \tag{15}$$

where

$$\begin{aligned} x_0 &= \frac{1}{4}\frac{\omega_2-\omega_1}{k_1\omega_2-k_2\omega_1}\ln(1-a)\ , \\ t_0 &= \frac{1}{4}\frac{k_2-k_1}{k_1\omega_2-k_2\omega_1}\ln(1-a)\ . \end{aligned} \tag{16}$$

The symmetry manifests itself in the log-log plots Fig.1, Fig.3, Fig.5, Fig.7, Fig.9 as an inversion symmetry with respect to the symmetry point

$$p = q = \sqrt{1-a}\ . \tag{17}$$

(iv) Although $p = 0$, $q = 0$ satisfies Eq.(11), in the first quadrant no curve starts from the origin. This is because for small p and q the first order terms dominate, and (for positive p and q) they are both negative.

(v) For $p \to \infty$ two asymptotes exist, namely, one at $q = 0$ and another one at $q = 1$. Similarly, for $q \to \infty$ we have asymptotes at $p = 0$ and $p = 1$. This result can easily be obtained since e.g. for large values of p the terms containing the highest (third) power of p dominate, i.e.,

$$k_2^3p^3q^2 - k_2^3p^3q = 0\ . \tag{18}$$

(vi) Near the above asymptotes the curve may be approximated by

$$q = \frac{k_1^3}{k_2^3}(1-a)\frac{1}{p} \quad \text{if} \quad p\to\infty\ ,\ q\to 0 \tag{19}$$

$$1-q = \frac{k_1^3}{k_2^3}\left[8 - a\left(4 + 6\frac{k_2}{k_1} - \left(\frac{k_2}{k_1}\right)^3\right)\right]\frac{1}{p} \quad \text{if} \quad p\to\infty\ ,\ q\to 1 \tag{20}$$

$$p = \frac{k_2^3}{k_1^3}(1-a)\frac{1}{q} \quad \text{if} \quad q\to\infty\ ,\ p\to 0 \tag{21}$$

$$1-p = \frac{k_2^3}{k_1^3}\left[8 - a\left(4 + 6\frac{k_1}{k_2} - \left(\frac{k_1}{k_2}\right)^3\right)\right]\frac{1}{q} \quad \text{if} \quad q\to\infty\ ,\ p\to 1 \tag{22}$$

This can be readily shown by taking into account the next-to-highest (second) power of the large variable.

The right hand sides of Eqs.(19), (21) are positive, thus the curve approaches the asymptote from above and from the right, respectively.§ The same is true for

§ Hence they lie in the first quadrant of the (p, q) coordinate system.

Eqs.(20), (22) in case of oppositely moving solitons ($\omega_1\omega_2 < 0$). In contrast, for solitons moving in the same direction ($\omega_1\omega_2 > 0$) the right hand sides of Eqs.(20), (22) can be both negative or positive, depending on the parameters k_1, k_2.

The above properties allow us to explain the possible topologies of the curve (11).



Figure 1. The zeros of Eq.(11) for solitons moving in the same direction. Left panel: Type I. case (at $k_1 = 1$, $k_2 = 1.5$), right panel: Type II. case (at $k_1 = 1$, $k_2 = 1.7$).

Under transformation (13) the point $p = 1$, $q = \infty$ goes over into $p = 1 - a$, $q = 0$, similarly, the point $p = \infty$, $q = 1$ goes over into $p = 0$, $q = 1 - a$. Finally, the point $p = 0$, $q = \infty$ goes over into $p = \infty$, $q = 0$. This allows only three possible topologies, according to the fact that the point $p = 1 - a$, $q = 0$ should be continuously connected with either $p = 0$, $q = 1 - a$ (Fig.1 left panel), or $p = \infty$, $q = 0$ (Fig.1 right panel), or $p = 1$, $q = \infty$. This last possibility is visualized again by the right panel of Fig.1 if p is exchanged with q, which, according to property 1 above, is equivalent with exchanging k_1 with k_2. Since these curves are intersections of a smooth surface with a plain, other possibilities are ruled out. We shall call the topology shown in the left panel of Fig.1 the Type I. case, and the topology shown in the right panel of Fig.1 the Type II. case. The topology obtained from the Type II. case via exchanging the axes will be called the Type $\overline{\text{II}}$. case. Note that the same transformation does not change the topology Type I.

While all the three situations do occur for solitons moving in the same direction, the Type I. case never occurs for oppositely moving solitons. This can be shown by considering the intersections of the curve (11) with the $p = q$ line, i.e., the zeros of

$$\xi \left[\xi^2 - (1-a)\right] \left[\xi^2 + b\xi + (1-a)\right] \ , \tag{23}$$

where $\xi = p = q$ and

$$b = 2 - a\frac{(k_1+k_2)^3}{k_1^3+k_2^3} \ . \tag{24}$$

Polynomial (23) always has zeros at $\xi = 0$ and $\xi = \pm\sqrt{1-a}$. The discriminant of the last quadratic factor is

$$D = b^2 - 4(1-a) = \frac{a(k_1+k_2)^6}{(k_1^3+k_2^3)^2}\left[a - \frac{12k_1k_2(k_1^3+k_2^3)}{(k_1+k_2)^5}\right] \ . \tag{25}$$

It is a simple exercise to show that the last factor on the right hand side of Eq.(25) is always negative in the case of oppositely moving solitons (cf. Eq.(4)). Hence, in that case only a single positive root exists, while the case Type I. shown in the left panel of Fig.1 requires three positive roots.

When changing the parameters k_1, k_2, the curves with different topologies go over into each other. If one considers solitons moving into the same direction, it is sufficient to consider the case $k_1 < k_2$, since the parameters must not coincide and the opposite case (i.e. $k_1 > k_2$) simply corresponds to the exchange of the axes. While the relative difference of parameters is sufficiently large, we have the topology Type II. If the relative difference is diminished, we get the topology Type I. The crossover between the two topologies if shown in left panel of Fig.2.



Figure 2. Crossover of the curves (11) having different topologies. Left panel: crossover from Type I. to Type II. for solitons moving in the same direction (at $k_1 = 1$, $k_2 = 1.57912575$), right panel: crossover from Type II. to Type $\overline{\text{II}}$. for solitons moving in opposite directions (at $k_1 = 1$, $k_2 = 1$).

One might wonder whether the middle branch of the curve touches both the other branches indeed at the same parameters. The answer is affirmative, and follows again from the symmetry (13). At the crossings of the branches one gets zero first derivatives in two independent directions‖, hence, one has simultaneously

$$f(p, q, k_1, k_2) = 0 \,, \tag{26}$$

$$\frac{\partial f(p, q, k_1, k_2)}{\partial p} = 0 \,, \tag{27}$$

$$\frac{\partial f(p, q, k_1, k_2)}{\partial q} = 0 \,, \tag{28}$$

where

$$f(p, q, k_1, k_2) = \sum_{n=1}^{5} \sum_{j=0}^{n} c_{nj} p^j q^{n-j} \,. \tag{29}$$

‖ This is actually an intersection of a saddle with the tangent plane.

Applying now the transformation (13) to a crossing point, it is straightforward to show that at the transformed point Eqs.(26)-(28) are also satisfied. Indeed, the two crossings go over into each other under transformation (13)¶.

In case of oppositely moving solitons, $k_1 = k_2$ is certainly possible, and a transition from Type II. to Type $\overline{\text{II}}$. occurs exactly when the two parameters coincide (cf. right panel of Fig.2).

3. Time evolution of the extrema

Extrema of the waves are the intersection points of Eqs.(11) and (12). As time goes on, the coefficient of the power function (12) changes from zero to infinity, and the corresponding curve$^+$ "sweeps through" the curve (11). According to the possible shapes of curve (11) we observe different scenarios, according to the number of maxima. As shown in Figs.3 and 4, in the Type I. case we always get three intersections, i.e., there are always two maxima and a minimum between them.



Figure 3. Type I. topology for intersections of the curve (11) and (12) at parameters $k_1 = 1.0$ and $k_2 = 1.5$ for solitons moving in the same direction. Different colors of the straight lines correspond to different time instants.

¶ The transformation is obviously an involution, i.e., it is equal to its own inverse.
$^+$ On the log-log plots the graph of Eq.(12) is a straight line.



Figure 4. Type I. topology as seen in wave pattern for solitons moving in the same direction. Parameters and time instants identified by the colors are the same as in Fig.3.

In contrast, in the Type II. and Type $\overline{\text{II}}$. cases there are periods when only a single maximum exists. However, one has to distinguish here two subcases:

a) If the middle branch is steep enough, namely, if

$$\frac{\partial \ln q}{\partial \ln p} > \frac{k_2}{k_1} \tag{30}$$

at the symmetry point (17), then at certain times the two maxima reappear (Figs.5 and 6). This situation will be called Type II.a (or Type $\overline{\text{II}}$.a).



Figure 5. Type II.a topology for intersections of the curve (11) and (12) at parameters $k_1 = 1.0$ and $k_2 = 1.6$ for solitons moving in the same direction. Different colors of the straight lines correspond to different time instants.



Figure 6. Type II.a topology as seen in wave pattern for solitons moving in the same direction. Parameters and time instants identified by the colors are the same as in Fig.5.

In such a case the initially well separated solitons (two maxima) coalesc, the remnant of the smaller soliton being only a "drooping shoulder" at the front side. Further on, the "shoulder" moves towards the maximum, it rises and develops a second maximum. Thus a shallow valley is created on the top of the wave. At later times these events take place in reversed order: the rear bank of the valley goes down, the corresponding maximum disappears and becomes a drooping shoulder at the rear side, then it develops a maximum again behind the taller wave and the two solitons are again separated.

b) If the steepness of middle branch is smaller than k_2/k_1, no valley is created on the top of the wave, as shown in Figs. 7 and 8. We shall call this case Type II.b (or Type $\overline{\text{II}}$.b).



Figure 7. Type II.b topology for intersections of the curve (11) and (12) at parameters $k_1 = 1.0$ and $k_2 = 1.8$ for solitons moving in the same direction. Different colors of the straight lines correspond to different time instants.



Figure 8. Type II.b topology as seen in wave pattern for solitons moving in the same direction. Parameters and time instants identified by the colors are the same as in Fig.7.

Both Type II.a and Type II.b can be observed in case of solitons moving in the same direction. In contrast, for oppositely moving solitons, only the case Type II.b (or Type $\overline{\text{II}}$.b) can exist. This can be proven on the basis of Eqs.(30), (17). Indeed, we have at the symmetry point

$$\frac{\partial \ln q}{\partial \ln p} = -\frac{4k_1^3(1-a) - a(k_2-k_1)^3 + \sqrt{1-a}(4k_1^3 - a(k_1+k_2)^3)}{4k_2^3(1-a) + a(k_2-k_1)^3 + \sqrt{1-a}(4k_2^3 - a(k_1+k_2)^3)}\,, \qquad (31)$$

which is smaller than k_2/k_1 for oppositely moving solitons, if $k_2 \geq k_1$.



Figure 9. Type II.b topology for intersections of the curve (11) and (12) at parameters $k_1 = 1.0$ and $k_2 = 1.5$ for solitons moving in opposite directions. Different colors of the straight lines correspond to different time instants.



Figure 10. Type II.b topology as seen in wave pattern for solitons moving in opposite directions. Parameters and time instants identified by the colors are the same as in Fig.9.

4. Parameter space

The scenarios described above are summarized in parameter space in Figs.11 and 12.



Figure 11. Scenarios shown in parameter space for solitons moving in the same direction. Legend: red, dark red: Type I. (see Fig.3 and Fig.4), white: Type II.a (see Fig.5 and Fig.6), gray: Type $\overline{\text{II}}$.a, green: Type II.b (see Fig.7 and Fig.8), dark green: Type $\overline{\text{II}}$.b.

For solitons moving in the same direction, at the border between Type II.a and Type II.b in parameter space (see Fig. 11.) the expression (31) is equal to k_2/k_1. This condition defines the border. As noted after Eq.(10), for both k_1 , $k_2 \ll 1$ and k_1 , $k_2 \gg 1$ the expression (31) depends only on the ratio k_2/k_1, hence the border looks linear. In fact, its slope slightly differs for small and large k values.

As discussed above, the border between between Type I. and Type II.a in parameter

space is given by Eqs.(26)-(29). Again, the border is not exactly a straight line, its slope is slightly different for large and small k values.

The shaded and the unshaded regions in Fig.11. are obtained by exchanging k_1 and k_2.

As noted before, the $k_1 = k_2$ line is not allowed.

For solitons moving in opposite directions, the parameter space is even simpler (see Fig.12.). All parameter values are allowed, and the crossover from Type II.b to Type $\overline{\text{II}}$.b occurs at $k_1 = k_2$.



Figure 12. Scenarios shown in parameter space for solitons moving in opposite directions. Legend: green: Type II.b (see Fig.9) and Fig.10), dark green: Type $\overline{\text{II}}$.b.

5. Nearly identical solitons

Let us consider now the situation when two nearly identical solitons interact. If they were strictly identical, we would obtain a single soliton solution rather than a two-soliton solution. Hence approaching the limit can be interesting. This situation corresponds to case I. Maxima correspond to intersections of the line (12) with the two outer segments of the graph of (11). Obviously, as time goes on, the initially well separated solitons approach each other, then, without coalescing, their distance grows again. The minimal distance between them may be estimated (cf. Fig.(13)) as

$$\delta x_{min} \geq -\frac{1}{k_1+k_2}\ln(1-a) = \frac{1}{k_1+k_2}\ln\left(\frac{(k_1+k_2)^2+\frac{1}{12}\left(\frac{\omega_1}{k_1}-\frac{\omega_2}{k_2}\right)^2}{(k_1-k_2)^2-\frac{1}{12}\left(\frac{\omega_1}{k_1}-\frac{\omega_2}{k_2}\right)^2}\right). \quad (32)$$

Evidently, δx_{min} diverges logarithmically as $k_1 \to k_2$.

If one considers now the corresponding waves (Fig.(14)), initially one sees two very similar solitons, the (slightly) taller one chasing the smaller one. When approaching, the tall soliton gradually loses its height and speed, at the same time, the smaller soliton gains hight and speed, the distance between them starts increasing, and eventually, we

see the original solitons again, but this time the smaller one chasing and the taller one escaping.



Figure 13. Intersections of the curve (11) and (12) at parameters $k_1 = 1.0$ and $k_2 = 1.1$ for solitons moving in the same direction.



Figure 14. Wave patterns at consecutive time instants for parameters $k_1 = 1.0$ and $k_2 = 1.1$. The solitons are moving in the same direction.

6. Summary and discussion

A simple classification scheme of the two soliton solutions of the Boussinesq equation have been presented. The scheme is based on the behavior of local maxima of the wave. We have shown that for solitons moving in the same direction there can be three different scenarios. In the Type I. case (see Figs. 3 and 4) there are two maxima all the time, separated by a minimum. In the Type IIa. case (see Figs. 5 and 6) initially, when the solitons are still separated, there are two maxima with a minimum in between. During

the collision the two solitons merge and only one maximum remains, the remnant of the other shows up only as a shoulder. Later on, however, the second maximum reappears and grows. Then the first maximum disappears for a while, but as the solitons become separated, it reappears and we have the initial solitons in a reversed ordering along the line. In the Type IIb. (see Figs. 7 and 8) case the separated solitons merge to a wave having a single maximum, and later on a second maximum reappears and the solitons separate again. For solitons moving in opposite directions only the Type IIb. case exists (see Figs. 9 and 10). In that case the wave numbers k_1 and k_2 may coincide (see Figs. 15 and 16). In contrast, for solitons moving in the same direction k_1 and k_2 must be different. If $k_1 \to k_2$ (see Figs. 13 and 14) we have an extreme Type I. case, namely, there remains a large minimal distance between the solitons all the time, and the chasing soliton loses in height, while the escaping soliton gains in height during the collision. As a result, eventually the two solitons change their ordering along the line, without any close contact.

Since our result are based on the long wave approximation, we expect that they should be observable in that limit. Also, the results for solitons moving in the same direction should follow from the Korteweg-de Vries equation as well, since that equation is obtained in the same approximation.



Figure 15. Intersections of the curve (11) and (12) at identical parameters $k_1 = k_2 = 1.0$ for solitons moving in opposite directions.



Figure 16. Wave patterns at consecutive time instants for identical parameters $k_1 = k_2 = 1.0$. The solitons are moving in opposite directions.

[1] G.F.Carrier and H.P.Greespan, *Water waves of finite amplitude on a sloping beach*, Fluid Mechanics, p.97, 1958.

[2] George F. Carrier, Tai Tei Wu and Harry Yeh, *Tsunami run-up and draw-down on a plane beach*, J. Fluid Mech. **475** (2003) 79 - 99.

[3] H.Bredmose et. al., *Experimental investigation and numerical modelling of steep forced water waves*, J. Fluid Mech. **490** (2003) 217.

[4] Kevin G. Lamb, *A numerical investigation of solitary internal waves with trapped cores formed via shoaling*, J. Fluid Mech. **451** (2002) 109.

[5] Kevin G. Lamb, *Shoaling solitary internal waves: on a criterion for the formation of waves with trapped cores*, J. Fluid Mech. **478** (2003) 81 - 100.

[6] A.R. Champneys, J.-M. Vanden-Droeck and G.J.Lord, *Do true elevation gravity-capillary solitary waves exist? A numerical investigation*, J. Fluid Mech. **454** (2002) 403 - 417.

[7] J.Kristian Sveen, Yakun Guo, Peter A. Davies and John Grue, *On the breaking of internal solitary waves at a ridge*, J. Fluid Mech. **469** (2002) 161 - 188.

[8] Ying Li and Frederic Raichlen, *Non-breaking and breaking solitary wave run-up*, J. Fluid Mech. **456** (2002) 295 - 318.

[9] Daohua Zhang and Allen T. Chwang, *On solitary waves forced by underwater moving objects*, J. Fluid Mech. **389** (1999) 119 - 135.

[10] Daohua Zhang and Allen T. Chwang, *Generation of solitary waves by forward- and backward-step bottom forcing*, J. Fluid Mech. **432** (2001) 341 - 350.

[11] R.S. Johnson, *Camassa-Holm, Korteweg-de Vries and related models for water waves*, J. Fluid Mech. **455** (2002) 63 - 82.

[12] Jiangbo Zhou, Lixin Tian, *Solitons, peakons and periodic cusp wave solutions for the Fornberg-Whitham equation*, arXiv:0908.0921v1

[13] N.Sugimoto, M.Masuda, K.Yamashita and H.Horimoto,*Verification of acoustic solitary waves*, J. Fluid Mech. **504** (2004) 271 - 299.

[14] F. Dias and J.-M. Vanden-Broeck, *Trapped waves between submerged obstacles*, J. Fluid Mech. **509** (2004) 93 - 102.

[15] Joseph B. Keller, *Shallow-water theory for arbitrary slopes of the bottom*, J. Fluid Mech. **489** (2003) 345 - 348.

[16] Atle Jensen, Geir K. Pedersen and Deborah J. Wood, *An experimental study of wave run-up at a steep beach*, J. Fluid Mech. **486** (2003) 161 - 188.

[17] Philip L.-F. Liu and Khaled Al-Banaa, *Solitary wave runup and force on a vertical barrier*, J. Fluid Mech. **505** (2004) 225 - 233.

[18] Alexander G. Voronovich, *Strong solitary internal waves in a 2.5-layer model*, J. Fluid Mech. **474** (2003) 85 - 94.

[19] Drazin P G and Johnson R S 2002 *Solitons: An Introduction* (Cambridge: Cambridge University Press)

[20] N. J. Zabusky, M. D. Kruskal Interaction of "Solitons" in a Collisionless Plasma and the Recurrence of Initial States. Phys. Rev. Lett. 15, 240 - 243 (1965)

[21] Clifford S. Gardner, John M. Greene, Martin D. Kruskal, and Robert M. Miura.
Method for Solving the Korteweg-de Vries Equation, Phys. Rev. Lett. 19, 1095 - 1097 (1967)

[22] Peter David Lax: Integrals of nonlinear equations of evolution and solitary waves, Comm. Pure Appl. Math. 21, 467 - 490(1968).

[23] M. J. Ablowitz and H. Segur: Asymptotic solutions of the Korteweg-de Vries equation, Stud. Appl. Math. 57, 13 - 44(1977).

[24] M. J. Ablowitz and P. A. Clarkson: Solitons, Nonlinear Evolution Equations and Inverse Scattering, volume 149 of London Mathematical Society Lecture Note Series Cambridge University Press, The Edinburgh Building, Shaftesbury Road, Cambridge, $CB2\ 2RU$, December (1991).

[25] S. P. Novikov, S. V. Manakov, L. P. Pitaevskii, and V. E. Zakharov: Theory of Solitons. The Inverse Scattering Method Plenum, New York(1984).

[26] Mark J. Ablowitz and Harvey Segur: Solitons and the Inverse Scattering Transform, volume 4 of SIAM Studies in Applied Mathematics SIAM Philadelphia(1981).
[27] Lei Wu, Jie-Fang Zhang and Lu Li, 2007 New J. Phys. **9** 69.
[28] S Mahmood, Mushtaq A and H Saleem 2003, New J. Phys. **5** 28.
[29] M. Tanaka, *The stability of solitary waves*, Phys. Fluids **29** (1986) 650.

[30] T. Kataoka and M. Tsutahara *Transverse instability of surface solitary waves*, J. Fluid Mech. **512** (2004) 211 - 221

[31] Kivshar Y S and Malomed B A 1989 *Rev.Mod.Phys.* **61** 765